

## Hazardous Effect Free From Decontamination

Hideharu Shintani\*

Faculty of Science and Engineering, Chuo University, 1-13-27, Kasuga, Bunkyo, 112-8551, Tokyo, Japan

\*Corresponding author: Hideharu Shintani, Faculty of Science and Engineering, Chuo University, 1-13-27, Kasuga, Bunkyo, 112-8551, Tokyo, Japan; TEL +81425922336; FAX +81425922336; E-mail: [shintani@mail.hinocatv.ne.jp](mailto:shintani@mail.hinocatv.ne.jp)

Received date: June 1, 2014, Accepted date: June 3, 2014, Published date: June 10, 2014


Copyright: © Fatme Al Anouti, 2014. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Editorial

Adventitious agents are those that are not inherent in the production of biopharmaceuticals. Microbial adventitious agents, Microbiocidal agents, help for inactivation of viruses, bacteria, fungi, mycoplasma, bacterial spore and sometimes prion. Bovine spongiform encephalopathy (BSE) agents, abnormal prions, are also potential adventitious agents. Adventitious agents can be introduced during establishment of cell lines, cell culture/fermentation, capture and downstream processing steps, formulation/filling, and even during drug delivery. Therapeutic biotechnology products must have excellent safety characteristics. However, the potential introduction of adventitious agents must continually be evaluated for safety. The testing procedures that is performed for this purpose is addressed in regulatory documents that include ICH guidelines, European U.S., and Japanese Pharmacopoeia (EP, USP and JP) documents. In some cases, 9 CFR (U.S. Code of Federal Regulations) and 21 CFR, 211 and 610 are also applicable.

Since biopharmaceuticals encompass many types of products, there is considerable variability in risks from adventitious agents. In all cases, however, the use of Good Manufacturing Practices (GMPs) (e.g. environmental controls, control of raw materials and personnel flow, and cleaning), suitable safety testing programs, and process validation (including evaluation for bacterial spore and sometimes prion

inactivation) helps to ensure patient confidence in biopharmaceuticals. As shown in the Table 1, prion is thought to be the most tolerable protein followed to bacterial spores. Therefore, prion and bacterial spore inactivation is the most important concern for avoiding deterioration. Some products have minimal inherent risk associated with the introduction of adventitious agents (e.g. recombinant products produced in bacteria). Other types of products, such as those used for cell or gene therapy, are often at the other end of the spectrum and may be associated with greater risk due to their inability to tolerate rigorous processing conditions. Greater potential risks are often associated with the use of human cells and animal-derived raw materials. Of particular concern are materials derived from bovine and porcine sources. When viable cells are a component of the product (as in ex vivo transduction), there may not be sufficient time to perform the required safety testing. In such cases, product may be released prior to completion of relevant tests for adventitious agents, although this testing is still mandated in order to demonstrate that the processes are being performed under adequate controls to maintain patient safety. A risk/benefit analysis determines whether the use of these products is warranted. We need to prevent and control of risks arising from various stages of production. We must confirm that several risks are associated with various sources and must not overlook the risks associated with several sources.

	Micro organism	Examples
	Prions	Scrapie, Creutzfeldt-jakob disease, Chronic wasting disease
	Bacterial Spores	<i>Bacillus</i> , <i>geobacillus</i> , <i>Clostridium</i>
	Protozoal oocysts	<i>cryptosporidium</i>
	Helminth eggs	<i>Ascaris</i> , <i>Enterobius</i>
	Mycobacteria	<i>Mycobacterium tuberculosis</i> , <i>M. terrae</i> , <i>M. Chelonae</i>
	Small Non enveloped viruses	<i>Polio virus</i> , <i>parvoviruses</i> , <i>papillomaviruses</i>
	Protozoal cysts	<i>Giardia</i> , <i>Acanthamoeba</i>
	Fungal spores	<i>Aspergillus</i> , <i>penicillium</i>
	Gram negative bacteria	<i>Pseudomonas</i> , <i>providencia</i> , <i>Escharichia</i>
	Vegetative fungi and algae	<i>Aspergillus</i> , <i>trichophyton</i> , <i>Candida</i> , <i>Chlamydomonas</i>
	Vegetative helminths and protozoa	<i>Ascaris</i> , <i>Cryptosporidium</i> , <i>Giardia</i>
	Large non enveloped viruses	<i>Adenovirus</i> , <i>rotaviruses</i>
	Gram positive bacteria	<i>Staphylococcus</i> , <i>Streptooccus</i> , <i>Enterococcus</i>

---

	Enveloped viruses	Human immunodeficiency virus, Hepatitis B virus, herpes simplex virus
--	-------------------	---

**Table 1:** General tolerance among microorganisms against inactivation agents. Prion is thought to be the greatest and next is spore. This order is unchanged. Citation is from McDonnell G.E. In Antisepsis, Disinfection, and Sterilization, ASM Press, Washington, 2007.