

## Changes in Physico-chemical Parameters at different Sites of Manasbal Lake of Kashmir, India

Gulzar Naik\*, Mudasir Rashid and Balkhi MH

Faculty of Fisheries, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Rangil, Ganderbal, Jammu and Kashmir, India

### Abstract

In an aquatic ecosystem, the density and diversity of organisms depends on availability and quality of water. During the present study an attempt was made to analyse some of the important physico-chemical parameters of the water samples at three different sites of the Manasbal Lake to check the pollution load. Various parameters were analysed for a period of six months on monthly intervals by following standard methods. Among the various parameters recorded the overall Air temperature ranged from 7.2°C to 30.2°C; Surface water temperature ranged from 6.1°C to 26.5°C; Secchi-disc transparency from 1.5 m to 4.5m; Conductivity varied from 149 to 292  $\mu\text{Scm}^{-1}$ ; Dissolved oxygen from 2.4 to 8.8  $\text{mg L}^{-1}$ ; Free  $\text{CO}_2$  from 2.0 to 14.0  $\text{mg L}^{-1}$ ; pH from 8.1 to 9.2; Total Alkalinity varied from 126 to 243  $\text{mg L}^{-1}$ ; Chloride values ranged from 5.9 to 33.5  $\text{mg L}^{-1}$ ; Total Hardness ranged from 119 to 218  $\text{mg L}^{-1}$ ; Ammonical Nitrogen varied from 27 to 193  $\mu\text{g L}^{-1}$ ; Nitrate Nitrogen varied from 31 to 286  $\mu\text{g L}^{-1}$ ; Ortho-Phosphate Phosphorous varied from 4.3 to 33.0  $\mu\text{g L}^{-1}$  and Phosphate Phosphorous ranged from 83.0 to 261  $\mu\text{g L}^{-1}$ . Almost all the above parameters showed that the pollution load is increasing especially at littoral sites due to agricultural run-off and human settlements disposing sewage, besides anthropogenic stresses in the catchment area. The pH of water of this lake was higher due to calcium intrusion from Kondabal area. Hence, the present study urges the need for immediate remedial measures for protection and conservation of this lake in order to save it from further deterioration.

**Keywords:** Manasbal lake; Physico-chemical parameters; Eutrophication; Pollution

### Introduction

The nature and health of any aquatic community is an expression of quality of the water. The quality of water affects the species composition, abundance, productivity and physiological conditions especially, the indigenous population of aquatic organisms [1]. The alteration in physico-chemical parameters leading to eutrophication has become widely recognized problem of water quality deterioration [2]. In recent years increase in human population, demand for food, land conversion and use of fertilizer have led to faster degradation of many freshwater resources [3,4]. The Kashmir valley located in the foothills of the Himalayas abounds in fresh water natural lakes that have also faced various anthropogenic activities and have resulted in heavy inflow of nutrients into these lakes from the catchment areas. These anthropogenic influences not only deteriorate the water quality but also affect the aquatic life in the lakes, as a result of which the process of aging of these lakes is hastened [5,6]. As a consequence, most of the lakes in the Kashmir valley are exhibiting eutrophication [7,8]. It is now quite common that most of the plain valley lakes of Kashmir are characterized by excessive growth of macrophytic vegetation, anoxic deep water layers and shallow marshy conditions along the peripheral regions and have high loads of nutrients in their waters [9]. The present paper includes study of physico-chemical parameters in water of Manasbal Lake, a tourist spot in Safapora town of District Ganderbal in Jammu and Kashmir.

### Materials and Methods

The Manasbal Lake is located at district Ganderbal about 32 kilometers away towards North West of Srinagar city in the State of Jammu and Kashmir, India. The lake is situated at an altitude of 1583 m asl and lies between 34°15' N latitude and 74°40' E longitude. The lake covers an area of 2.81  $\text{km}^2$  with maximum depth of 13 m. The volume of water has been estimated as  $12.8 \times 10^6 \text{ m}^3$  [10]. The lake catchment covers an area of about 22  $\text{km}^2$ . The lake has no major inflows and its

water supply is chiefly derived from internal springs and precipitation. The lake is surrounded by many villages among which Kondabal and Hanji Mohalla are very important as far as health and vigour of the lake is concerned. Kondabal is the place of kilns, quarries and mines is significant as far as the calcium intrusion into the lake is concerned. Hanji Mohalla is inhabited by fishers from which lot of organic matter and other pollutants enter into the lake. Manasbal is a marl lake, has predominantly rural surrounding. Currently, one of the visible problems within the lake is eutrophication.

### Study sites (Figure1)

Three different sites were selected for the present study on the basis of pollution load as:

**Site-I:** Littoral site near Manasbal Garden (Ghat) and Kondabal and usually with high density of aquatic macrophytes.

**Site-II:** Central limnetic site representing the open water zone with a maximum depth and relatively a lesser density of macrophytic vegetation, the water at this site is relatively clear with a greater degree of visibility.

**Site-III:** Littoral site near Hanji Mohalla which is surrounded by residential hamlets, exposed to myriad of interference, such as emission of domestic sewage and washing of clothes and other activities.

\*Corresponding author: Gulzar Naik, Faculty of Fisheries, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Rangil, Ganderbal, Jammu and Kashmir, 190006, India, Tel: 0194 246 2159; E-mail: [mudasir.rathero@gmail.com](mailto:mudasir.rathero@gmail.com)

Received July 03, 2015; Accepted November 04, 2015; Published November 24, 2015

**Citation:** Naik G, Rashid M, Balkhi MH (2015) Changes in Physico-chemical Parameters at different Sites of Manasbal Lake of Kashmir, India. Fish Aquac J 6: 148. doi:10.4172/2150-3508.1000148

**Copyright:** © 2015 Naik G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Map of Manasbal Lake showing different study sites.

### Sampling

Water samples were collected from the selected sites on monthly basis for a period of six months from July 2010 to December 2010. The water samples were collected during second week of each month from different study sites. Samples were analysed according to standard methods as described by Welch, Golterman and Clymo and APHA [11-13]. Some of the parameters like temperature, transparency, pH were determined at the field site. For dissolved oxygen, water samples were fixed on the spot in BOD bottles as per the Winkler’s modified method where as the other parameters were analysed at the laboratory.

### Results and Discussion

The maximum and minimum values of various physico-chemical parameters at three different sites is given in (Table 1).

#### Temperature

In the present study, air temperature ranged between 7.2°C in December at site-II to a maximum of 30.2°C in July at site-I. The lowest water temperature observed in December at site-II was 6.1°C and highest was observed in July at site-I viz., 26.5°C. This is because of the shallowness of the lake and consequently the volume of water in contact with air. Therefore, a close relationship was recorded between water temperature and air temperature which is in line with the findings of Zutshi [14].

#### Transparency

Transparency is one of the important physical properties of water

and is indicative of degree to which sunlight can pass through the water. During the present study the values for transparency obtained at site-II ranged between 4.5 m (December) to 3.3 m (July) and at site-III ranged between 2.4 m (December) to 1.5 m (July). However, on the whole Secchi disc transparency was low at the littoral sites being minimum at site-III which may be due to the increased suspended matter, silt load and humus brought in from the catchment area as well as due to rich macro-vegetation. Transparency was higher in colder and lower in warmer months and this condition has been attributed to different factors viz., plankton population, settling of materials in calm weather, suspension of phytoplankton in water, glacial silt e.t.c [14-17].

#### Conductivity

The conductivity value is an indicator of the total nutrient level of a water body. Using specific conductivity as an index, values higher than 200 µS/cm show higher level of nutrient enrichment [18]. During the present study conductivity reported was within the range of 149 µS/cm in December at site-II to 292 µS/cm in August at site- III. The values were higher at site-III because of more polluted water. A high level of conductivity reflects the pollution status as well as trophic level of the lake [19].

#### Dissolved oxygen

Dissolved oxygen is an important regulator of metabolic processes of the organisms and also the community as a whole [20]. Its concentrations in natural water depend on the physical, chemical and biological activities in the water body [21]. During the present study the highest dissolved oxygen level was observed at site-II in December viz., 8.8 mg/l and lowest in July at site-III viz., 2.4 mg/l. The higher values of dissolved oxygen in colder and lower values during warmer months, clearly reflecting an inverse relation of dissolved oxygen with the temperature. Dissolved oxygen levels during winter can also be attributed to low biological activity [22].

#### Free carbon dioxide

Free carbon dioxide of most waters is not present in large quantities because of its reaction in the carbonate equilibrium and exchange with atmosphere [1]. In the present study lowest Co<sub>2</sub> concentration was observed during July at site-I viz., 2 mg/l where as the highest concentration of Co<sub>2</sub> was reported at site-III viz., 14 mg/l in December. Slight temperature variation can alter solubility of oxygen and carbon dioxide in water [23]. The higher values towards winter may be due to

S.no	PARAMETER	SITE-I		SITE-II		SITE-III	
		MAX	MIN	MAX	MIN	MAX	MIN
1.	AIR TEMPERATURE (°C)	30.2 (JULY)	7.9 (DEC.)	29.0 (JULY)	7.2 (DEC.)	29.8 (JULY)	7.5 (DEC.)
2.	WATER TEMPERATURE (°C)	26.5 (JULY)	6.6 (DEC.)	25.8 (JULY)	6.1 (DEC.)	26.1 (JULY)	6.3 (DEC.)
3.	SECCHI TRANSPARENCY (m)	2.9 (DEC.)	1.8 (JULY)	4.5 (DEC.)	3.3 (JULY)	2.4 (DEC.)	1.5 (JULY)
4.	CONDUCTIVITY (µS/cm)	280 (AUG.)	192 (DEC.)	170 (AUG.)	149 (DEC.)	292 (AUG.)	268 (DEC.)
5.	DISSOLVED OXYGEN (mgL <sup>-1</sup> )	8.0(DEC.)	4.4 (JULY)	8.8(DEC.)	5.2 (JULY)	7.2(DEC.)	2.4 (JULY)
6.	FREE CARBONDIOXIDE (mgL <sup>-1</sup> )	12.0(DEC.)	2.0(JULY)	10.0(DEC.)	2.0(JULY)	14(DEC.)	4.0 (JULY)
7.	P <sup>H</sup>	9.2 (JULY)	8.5 (DEC.)	8.9 (AUG.)	8.2(DEC.)	8.9 (AUG.)	8.1 (DEC.)
8.	TOTAL ALKALINITY (mgL <sup>-1</sup> )	237 (DEC.)	175 (JULY)	154 (DEC.)	126 (JULY)	219 (DEC.)	163 (JULY)
9.	TOTAL HARDNESS (mgL <sup>-1</sup> )	218 (SEPT.)	161 (NOV.)	137 (SEPT.)	119(NOV.)	178(OCT.)	146 (DEC.)
10.	CHLORIDE (mgL <sup>-1</sup> )	21.2 (DEC.)	8.8 (JULY)	12.3 (DEC.)	5.9 (JULY)	33.5 (DEC.)	18.0 (AUG.)
11.	PHOSPHATE PHOSPHOROUS (µg <sup>-1</sup> )	202 (JULY)	143 (DEC.)	142 (JULY)	83 (DEC.)	261 (JULY)	189 (DEC.)
12.	ORTHO-PHOSPHATE PHOSPHOROUS (µg <sup>-1</sup> )	29 (OCT.)	12 (DEC.)	17 (AUG.)	4.3 (DEC.)	35 (OCT.)	17 (DEC.)
13.	AMMONICAL NITROGEN (µg <sup>-1</sup> )	138 (DEC.)	84 (JULY)	63 (DEC.)	27 (JULY)	193 (DEC.)	109 (JULY)
14.	NITRATE NITROGEN (µg <sup>-1</sup> )	201 (JULY)	116 (OCT.)	109 (JULY)	31 (NOV.)	286 (JULY)	178 (NOV.)

Table 1: The maximum and minimum values of various physico-chemical parameters at three different sites.

the decreasing temperature and addition of high organic matter from catchment area.

### Hydrogen ion concentration (pH)

All chemical and biological reactions are directly dependent upon pH of the water system [24]. The pH of this lake was found higher in all months due to addition of calcium in water from Kondabal. The highest value of pH was observed during July at site-I viz., 9.2 where as the lowest value was observed at site-III in December viz., 8.1. Fluctuations in pH seemed to be influenced by the changes in the rate of photosynthesis and decomposition and this was in conformity with the conclusion drawn by Zutshi and Khan [25]. Generally lower pH values were recorded in cold months.

### Total alkalinity

The total alkalinity in the lake was less in July at site-II viz., 126 mg/l and highest at site-I viz., 237 mg/l in December. The increase in alkalinity during cold months may be due to the accumulation of bicarbonate ions as the rate of their uptake is declined and the decrease in warmer months is due to the decrease in bicarbonate ions because of its utilization by luxuriant growth of phytoplankton and macrophytes. These findings were in agreement with Sahai and Shrivastava [26]. Balkhi observed that alkalinity of water is mainly due to bicarbonates of calcium and magnesium [27]. Total alkalinity has been used as a rough index of lake productivity [28]. Moyle designated the lake water having alkalinity values upto 40 mg/l as soft, those with 40-90 mg/l as medium hard and those with values above 90 mg/l as hard type [29]. During present study alkalinity was reported above 126 mg/l at all the sites of the lake.

### Total hardness

The two most common bivalent cations of lake water are Ca and Mg, usually account for most of the hardness. The carbonate hardness of water which includes the portion equivalent to carbonate and bicarbonate in water has been called temporary hardness. Permanent hardness is caused by sulphates and chlorides of bivalent cations. During the present study highest total hardness value was observed during September at site-I viz., 218 mg/l and lowest was observed in November viz., 119 mg/l at site-II. Water with hardness below 60 mg/l is soft, between 61 to 120 mg/l is moderately hard, 121 to 180 mg/l is hard and more than 181 mg/l is very hard [35].

### Chloride content

Chloride content in water is regarded as an indication of organic load of animal origin from the catchment area [30]. High chloride concentration in water body is related to organic pollution of animal origin [31]. During the present study highest chloride concentration was observed at site-III viz., 33.5 mg/l in November and lowest in July viz., 5.9 mg/l at site-II. The higher values at site-III may be due to the activities like washing of clothes, drying of dung cakes at the catchment area, domestic wastes etc.

### Phosphorous and ortho-phosphorous

Phosphorus plays a central role in eutrophication of water bodies. The low orthophosphate-phosphorous content in waters is due to the formation of an insoluble calcium-phosphate complex. Such a phenomenon functions as scavenger of some inorganic nutrients and also acts as a removal agent of dissolved organic matter by absorption. During the present investigation highest concentration of orthophosphate-phosphorous was found at site-III in October

viz., 33 µg/l and lowest at site-II in December viz., 4.3 µg/l. Highest concentration of phosphate-phosphorus was found at site-III in July viz., 261 µg/l and lowest was found at site-II in December viz., 83 µg/l. This can be attributed to the fact that the major sources of phosphorus are domestic sewage, detergents and agricultural effluents. Hutchinson related the increased phosphorus concentration to sewage contamination [20]. Correll is of the opinion that the formation of algal bloom is due to high phosphorus concentration, which leads to the eutrophication as a result of which several water bodies have perished [32].

### Ammonical-nitrogen

Ammonia is basically a product of decomposition. Its concentration during the study period was generally low because of the utilization of nitrogenous compounds like NH<sub>4</sub>-N by plants. During present study highest concentration of ammonical nitrogen was observed in December at site-III viz., 193 µg/l and lowest at site-II in July viz., 27 µg/l. High concentration at site-III may be due to the entry of domestic sewage and use of other nitrogenous pollutants like fertilizers in the catchment area. The low value of ammonical nitrogen during warmer months is attributed to the photosynthetic assimilation by autotrophs during their growth in summer [33].

### Nitrate-nitrogen

The presence of nitrates in any aquatic ecosystem depends on the activity of nitrifying bacteria. The conversion of nitrates from ammonia in nitrification process by *Nitrobacter* and *Nitrosomonas* chiefly depends upon the presence of oxygen. Nitrate is the common form of inorganic nitrogen entering freshwater from the draining basin, ground water and precipitation and mostly occurs in low concentration [34]. During the present study highest concentration of nitrate was observed in July at site-III viz., 286 µg/l and lowest at site-II in November (31 µg/l). This can be attributed to its locking up in luxuriant macrophytic population.

### Conclusion

From the present study it can be inferred that the littoral sites were more polluted and eutrophic than central limnetic site. The most polluted area of lake was site-III (Hanji Mohalla) followed by site-I near Manasbal Garden (Ghat) and the least polluted was site-II (Central site). The higher values of nitrogen, phosphorus, chloride, alkalinity, hardness, conductivity, free CO<sub>2</sub> and lower values of dissolved oxygen and transparency are clearly pointing towards higher trophic status of the lake especially at littoral sites. Some of the possible causes of deterioration of water quality of the lake include agricultural runoff and human settlements disposing sewage, besides anthropogenic stresses in the catchment. Hence, the present study urges the need for immediate remedial measures for protection and conservation of this lake in order to save it from further deterioration.

### References

1. Wetzel RG (2001) Limnology: Lake and river ecosystems. Academic Press. NY. USA, pp: 1006.
2. Kim B, Park J, Hwang G, Jun M, Choi K (2001) Limnology 2: 223-229.
3. Ray D, RavindarRao R, Bhoi AV, Biswas AK, Ganguly AK (1999) Environ. Monit. Assess 61: 387-398.
4. Carpenter RS (2005) PNSA Pp: 1-4.
5. Odada EO, Olago DO, Kuliindwa K, Ntiba M, Wandiga S, et al. (2004) Mitigation of Environmental Problems in Lake Victoria, East Africa: Causal Chain and Policy Options Analyses. Ambio 33: 13-23.

6. Li R, Dong M, Zhao Y, Zhang L, Cui Q, et al. (2007) Assessment of Water Quality and Identification of Pollution Sources of Plateau Lakes in Yunnan (China). *Journal of Environmental Quality* 36: 291-297.
7. Kaul V (1979) Water Characteristics of Some Fresh Water Bodies of Kashmir. *Current Trends in Life Sciences* 9: 221-246.
8. Khan MA (2008) Chemical Environment and Nutrient Fluxes in a Flood Plain Wetland Ecosystem, Kashmir Himalayas, India. *Indian Forester*. 134: 505-514.
9. Jeelani G, Shah AQ (2006) Geochemical characteristics of Water and Sediment from the Dal Lake, Kashmir Himalaya: Constraints on Weathering and Anthropogenic activity. *Environmental Geology* 50: 112-123.
10. Yousuf AR (1992) Biotic Communities and their role in the tropic conditions in Kashmir Himalayan Lakes, Technical Report submitted by the University of Kashmir, CSIR Research Project.
11. Welch PS (1948) *Limnology*. Mcgraw Hill Publications.
12. Golterman HL, Clymo RS (1969) *Methods for Chemical Analysis of Freshwater*. I.B.P. Handbook No : 8. Blackwell Scientific Publications, Oxford.
13. American Public Health Association (2005) *Standard methods for examination of water and wastewater*. APHA, AWWA, WPCF, Washington DC, USA.
14. Zutshi DP, Subla BA, Khan MA, Wangane A (1980) Comparative limnology of nine lakes of Jammu and Kashmir Himalayas. *Hydrobiologia* 80: 101-112.
15. Zutshi DP, Vass KK (1970) High altitude lakes of Kashmir, *Ichthyologia* 10: 12-15.
16. Spurr B (1975) *Limnology of Bird pond, Ross Island, Antarctica*, NZJ Mar Freshwat 9: 547-562.
17. Zutshi DP, Khan MA (1978) On lake typology of Kashmir. *Environ, Physiol. Ecol. Plants* pp: 465-472.
18. Rawson DS (1960) A Limnological comparison of twelve large lakes in Northern Saskatchewan. *Limnol. Oceanogr* 5: 195-211.
19. Shashtree NK, Islam MS, Pathak S, Afshan M (1991) Studies on the Physico-chemical dimensions of the lentic hydrosphere of RavindraSarover (Gaya). In *current trends in Limnology*. (Nalin K, Shashtreed). Narender Pub. House, N. Delhi pp: 132-152.
20. Hutchinson GE (1967) *A Treatise on Limnology*. John Wiley and Sons, NY Indian J. Ecol 25: 83-92.
21. Zutshi DP, Vass KK (1978) Variations in the water quality of some Kashmir Lakes *Trop. Ecol* 14: 182-196.
22. Qadri MY, Naqash SA, Shah GM, Yousuf AR (1981) Limnology of two trout streams of Kashmir. *J. Inst. Sci* 63: 137-147.
23. Wetzel RG (1975) *Limnology*. Mc-Graw Hill Book Co. Inc, New York pp: 538.
24. Sreenivasan A (1967) Application of limnological and primary productivity studies in fish culture. *F.A.O.F. sh Report* pp: 44: 101-103.
25. Zutshi DP, Khan MA (1988) Eutrophic gradient in the Dal lake, Kashmir. *Indian J. Env. Health* 30: 4.
26. Sahai R, Srivastava VC (1976) Physico-chemical complexes and its relationship with macrophytes of Chilka Lake. *Geobios* 3: 15-19.
27. Balkhi MH, Yousuf AR, Qadri MY (1987) *Hydrobiology of Anchar Lake, Kashmir*. *Comp. Physiol. Ecol* 12: 131-139.
28. Moyle JB (1956) Relationship between the chemistry of Minnesota surface waters and wild life management. *J Wild Mgt* 20: 303-320.
29. Moyle JB (1945) Some chemical factors influencing the distribution of aquatic plants in Glubokoe in relation to predation and composition. *Hydrobiol.* 141: 33-43.
30. Kumar A, Mandal KK, Kumar A (2004) Aspects of water pollution scenario in lentic and lotic freshwater namely pond and river in Jharkhand, India. *Water pollution by Prof. Arvind Kumar, A.P.H. Publishing Corp, S Ansari Road, Daryaganj, New Delhi* pp: 1-24.
31. Thresh JC, Suckling EV, Beale JE (1944) *The Examination of water and water supplies*. (EW Taylor) London.
32. Correl DL (1998) The role of phosphorus in the eutrophication of receiving waters, a review. *J Environ Quality (USA)* 27: 261-266.
33. Pandit AK (1999) Trophic structure of plankton community in some typical wetlands of Kashmir, India. *Limnological Research in India (SR Mishra) Daya Publishing House, Delhi*, pp: 190-224.
34. Wetzel RG (1983) *Limnology*. Saunders college of Business, New York.
35. Durfor CN, Becker E (1964) *Survey Water Quality Paper 1812: 1-364*.

**Citation:** Naik G, Rashid M, Balkhi MH (2015) Changes in Physico-chemical Parameters at different Sites of Manasbal Lake of Kashmir, India. *Fish Aquac J* 6: 148. doi:[10.4172/2150-3508.1000148](https://doi.org/10.4172/2150-3508.1000148)

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>